

(PCT Rule 61.2)

To:

Commissioner
US Department of Commerce
United States Patent and Trademark
Office, PCT
2011 South Clark Place Room
CP2/5C24
Arlington, VA 22202
ETATS-UNIS D'AMERIQUE
in its capacity as elected Office

Date of mailing (day/month/year) 26 January 2001 (26.01.01)	ETATS-UNIS D'AMERIQUE in its capacity as elected Office
International application No. PCT/IB00/00542	Applicant's or agent's file reference 023406.00003
International filing date (day/month/year) 27 April 2000 (27.04.00)	Priority date (day/month/year) 27 April 1999 (27.04.99)
Applicant HAMUNEN, Antti	

1. The designated Office is hereby notified of its election made:

☒ in the demand filed with the International Preliminary Examining Authority on:

27 November 2000 (27.11.00)

☐ in a notice effecting later election filed with the International Bureau on:

2. The election ☒ was
☐ was not

made before the expiration of 19 months from the priority date or, where Rule 32 applies, within the time limit under Rule 32.2(b).

<p>The International Bureau of WIPO 34, chemin des Colombettes 1211 Geneva 20, Switzerland</p> <p>Facsimile No.: (41-22) 740.14.35</p>	<p>Authorized officer</p> <p>S. Mafla</p> <p>Telephone No.: (41-22) 338.83.38</p>
---	--

INTERNATIONAL SEARCH REPORT

In International Application No

PC 00/00542

A. CLASSIFICATION OF SUBJECT MATTER
IPC 7 C11B13/00

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 7 C11B C11D C07J C09F

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the International search (name of data base and, where practical, search terms used)

WPI Data, EPO-Internal, PAJ, CHEM ABS Data, PIRA

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	W0 96 10033 A (UNIV BRITISH COLUMBIA) 4 April 1996 (1996-04-04) page 9, line 16 -page 10, line 10; example 1	1-8, 17-25, 34-47
A	DATABASE CAPLUS 'Online! CHEMICAL ABSTRACTS; AN 1979:493367, XP002143291 abstract & FI 55 679 C (LINOTEK OY) 22 September 1978 (1978-09-22) cited in the application -/--	1,3-9, 17,18, 20-26,39

☒ Further documents are listed in the continuation of box C.☒ Patent family members are listed in annex.

* Special categories of cited documents :

- *A* document defining the general state of the art which is not considered to be of particular relevance
- *E* earlier document but published on or after the international filing date
- *L* document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
- *O* document referring to an oral disclosure, use, exhibition or other means
- *P* document published prior to the international filing date but later than the priority date claimed

- *T* later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
- *X* document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
- *Y* document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art
- *&* document member of the same patent family

Date of the actual completion of the international search

24 July 2000

Date of mailing of the international search report

07/08/2000

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2
NL - 2280 HV Rijswijk
Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,
Fax: (+31-70) 340-3016

Authorized officer

Saunders, T

INTERNATIONAL SEARCH REPORT

International Application No

IB 00/00542

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	US 3 965 085 A (HOLMBOM BJARNE ET AL) 22 June 1976 (1976-06-22) cited in the application examples 3,5	1-8, 17-26, 33-36
A	EP 0 122 322 A (UNION CAMP CORP) 24 October 1984 (1984-10-24) claims	1,3-8, 17,18, 20-25

INTERNATIONAL SEARCH REPORT

Information on patent family members

International Application No

PCT/JP 00/00542

Patent document cited in search report		Publication date	Patent family member(s)	Publication date
WO 9610033	A	04-04-1996	AU 3559795 A	19-04-1996
			BG 101354 A	31-03-1998
			BR 9509119 A	28-10-1997
			CA 2201328 A	04-04-1996
			CN 1166174 A	26-11-1997
			CZ 9700927 A	18-02-1998
			EP 0783514 A	16-07-1997
			HU 77522 A	28-05-1998
			JP 10506394 T	23-06-1998
			LT 97074 A, B	27-10-1997
			LV 11900 A	20-12-1997
			LV 11900 B	20-05-1998
			MD 970178 A	28-02-1999
			NO 971304 A	14-05-1997
			PL 319384 A	04-08-1997
			SK 41597 A	05-11-1997
			US 5770749 A	23-06-1998
FI 55679	C	22-09-1978	FI 770873 A, B,	22-09-1978
US 3965085	A	22-06-1976	CA 1021221 A	22-11-1977
			JP 50038705 A	10-04-1975
			SE 417107 B	23-02-1981
			SE 7407901 A	30-12-1974
EP 0122322	A	24-10-1984	US 4422966 A	27-12-1983
			AT 25400 T	15-02-1987
			CA 1202302 A	25-03-1986
			FI 834818 A, B,	19-09-1984

PCTINTELLECTUAL PROPERTY ORGANIZATION
International Bureau

INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁷ : C11B 13/00	A1	(11) International Publication Number: WO 00/65004 (43) International Publication Date: 2 November 2000 (02.11.00)
(21) International Application Number: PCT/IB00/00542 (22) International Filing Date: 27 April 2000 (27.04.00) (30) Priority Data: 60/131,304 27 April 1999 (27.04.99) US (71) Applicant (for all designated States except US): STEROL TECHNOLOGIES LTD. [FI/FI]; P.O. Box 101, FIN-21201 Raisio (FI). (72) Inventor; and (75) Inventor/Applicant (for US only): HAMUNEN, Antti [FI/FI]; Lumparlankatu 13 C 20, FIN-21200 Raisio (FI).		(81) Designated States: AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CR, CU, CZ, DE, DK, DM, DZ, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, TZ, UA, UG, US, UZ, VN, YU, ZA, ZW, ARIPO patent (GH, GM, KE, LS, MW, SD, SL, SZ, TZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG). Published <i>With international search report.</i>
(54) Title: EXTRACTION AND ISOLATION METHOD (57) Abstract A method for separating neutral substances from a soap containing the neutral substances, the method including the steps of: providing a mixture containing (1) the soap containing the neutral substances, (2) water optionally containing sodium sulfate and (3) a C ₁ -C ₁₀ hydrocarbon solvent, and heating the mixture to a temperature of at least 140 °C, to obtain a soap phase and a hydrocarbon phase containing the neutral substances, the heating step being conducted in a closed system under pressure, wherein the pressure in the system is at least equal to the vapor pressure of the mixture at the temperature used in the heating step; separating the hydrocarbon phase from the soap phase; and optionally separating the neutral substances from the hydrocarbon phase.		

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav	TM	Turkmenistan
BF	Burkina Faso	GR	Greece		Republic of Macedonia	TR	Turkey
BG	Bulgaria	HU	Hungary	ML	Mali	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MN	Mongolia	UA	Ukraine
BR	Brazil	IL	Israel	MR	Mauritania	UG	Uganda
BY	Belarus	IS	Iceland	MW	Malawi	US	United States of America
CA	Canada	IT	Italy	MX	Mexico	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NE	Niger	VN	Viet Nam
CG	Congo	KE	Kenya	NL	Netherlands	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NO	Norway	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's	NZ	New Zealand		
CM	Cameroon		Republic of Korea	PL	Poland		
CN	China	KR	Republic of Korea	PT	Portugal		
CU	Cuba	KZ	Kazakstan	RO	Romania		
CZ	Czech Republic	LC	Saint Lucia	RU	Russian Federation		
DE	Germany	LI	Liechtenstein	SD	Sudan		
DK	Denmark	LK	Sri Lanka	SE	Sweden		
EE	Estonia	LR	Liberia	SG	Singapore		

EXTRACTION AND ISOLATION METHOD

This application claims priority under 35 USC § 119(e) of U.S. Provisional Application Serial No. 60/131,304, which is hereby incorporated by reference.

5

Field of the Invention

The present invention relates to a method for the separation of useful substances from soaps, such as pulping soap obtained as a by-product in the production of sulfate cellulose or soaps obtained from deodorized distillates of plant oils. More precisely the invention relates to a method for extracting
10 unsaponifiable neutral substances from different kind of soaps. The invention also relates to a method for extracting, separating and purifying sterols from soaps.

Background of the Invention

Soaps contain a lot of alkali salts of fatty and/or resin acids and variable
15 amounts of unsaponifiable neutral substances consisting of e.g. phytosterols, terpene alcohols, fatty alcohols, and aldehydes, ethers, ketones as well as hydrocarbons. Small amounts of fibres, lignins and inorganic salts also exist as impurities in the soap. The amount of neutral substances varies depending on the wood material used for pulping.

20 Tall oil, made from soaps by acidulation is a valuable raw material for the chemical industry, but generally the neutral substances included in it are not utilized at all. However, from the neutral substances many valuable raw materials can be isolated, such as phytosterols, which can be used e.g. in the food, pharmaceutical and cosmetic industry. Because of the many beneficial
25 characteristics of the phytosterols, the need thereof has grown enormously during the last years. Besides producing hormone preparations by steroid transformations they are used in foodstuffs or medical preparations especially for

lowering serum total and LDL cholesterol levels. Most preferable for this kind of use are the stanols and especially the stanol esters, which can be produced e.g. by hydrogenation and esterification of phytosterols.

One method for extracting neutral substances from soaps was proposed
5 in US 2,866,781. In the method proposed therein, esters of a relatively low molecular weight aliphatic acid are used as extraction solvents for aqueous soap solutions. These extraction solvents are, however, not very selective for the neutral substances.

US 3,965,085 discloses a method for refining soaps in which the
10 unsaponifiable neutral substances included in the soap are separated by means of extraction from an aqueous solution containing low molecular weight ketones, such as acetone or methyl ethyl ketone, using as the water-immiscible solvent an aliphatic, aromatic, alicyclic or halogenated hydrocarbon. In the Examples of this patent, extraction is conducted at temperatures below 75°C. The added
15 ketone is essential in the disclosed extraction method, since it prevents the formation of emulsions in the soap solution and thereby facilitates the extraction process.

US 2,530,809 discloses the extraction of neutral substances from tall oil
soaps using a mixture of the soaps with a lower alcohol and water. Suggested
20 as solvents for the unsaponifiable material are alkanes or other petroleum fractions, ethers, aromatic hydrocarbons or chlorinated hydrocarbons. The extraction procedure is conducted at temperatures of between 80 and 120°F (which is between about 27 and 49°C). In this case, the lower alcohol acts as an emulsion breaker.

25 Problems with the two extraction methods described in US 3,965,085 and US 2,530,809 are particularly focused on their use of a two-component solvent system, which involves difficult solvent regeneration systems in large scale production. Complete removal of the alcohol (US '809) or the ketone (US '085)

from the soap solution is also technically impossible, because of foaming problems. The remaining solvent is also to some extent harmful for further processing of the soap. If present, it causes problems e.g. in the distillation of crude tall oil. But, above all, the inevitable solvent losses in these methods make
5 them not very economic and these methods do not take into account environmental aspects.

FI 55 679 discloses an extraction method wherein a long chain C_{12} - C_{35} hydrocarbon in combination with water is used for extracting neutral substances from soap. This method suffers from the drawbacks involved in using long
10 chain hydrocarbons, which need very high temperatures for regeneration and therefore are not particularly suitable in large scale production. Another problem involved with this method is that the long chain hydrocarbons and the light fractions of the neutral substances have boiling points close to each other, and the recirculation of this solvent may therefore lead to higher amounts of
15 light fractions of the neutral substances in the process system.

All the prior disclosed methods for extracting neutral substances from soaps have some technical problems, which have been solved by the methods according to the present invention.

Summary of the Invention

20 It has now surprisingly been realized that neutral substances can be extracted from different kind of soaps by the use of a short chain (C_1 - C_{10}) hydrocarbon solvent and an aqueous solution of the soaps. This allows the best regeneration system for the solvent, and the problems associated with the use of long chain hydrocarbons are avoided. No other organic solvent is needed in
25 the method of the invention because, instead of using additives such as alcohols or ketones as in the art, emulsion breaking can be accomplished by increasing

the temperature of the mixture to at least 140°C, and thereby the phase separation needed for the extraction to proceed is accomplished.

In a first aspect of the present invention there is provided a method for the separation of neutral substances from soaps comprising the steps of:

- 5 extracting neutral substances into a hydrocarbon phase by mixing the soap with water optionally containing sodium sulfate, and with a short chain hydrocarbon solvent at ambient or elevated temperature, and raising the temperature of the mixture to at least 140°C, to obtain a hydrocarbon phase containing neutral substances, and a soap phase, the extraction step being
10 conducted in a closed system, wherein the pressure in the system is at least equal to the vapor pressure of the extraction mixture at the temperature used for extraction,

separating the hydrocarbon phase containing neutral substances from the soap phase, and

- 15 optionally separating the neutral substances from the hydrocarbon phase.

In a second aspect of the present invention there is provided a method for the separation of sterols from soaps comprising the steps of:

- extracting neutral substances into a hydrocarbon phase by mixing the soap with water optionally containing sodium sulfate, and with a short chain (C₁-C₁₀)
20 hydrocarbon solvent at ambient or elevated temperature, and raising the temperature of the mixture to at least 140°C, to obtain a hydrocarbon phase containing neutral substances, and a soap phase, the extraction step being conducted in a closed system, wherein the pressure in the system is at least equal to the vapor pressure of the extraction mixture at the temperature used for
25 extraction,

separating the hydrocarbon phase containing neutral substances from the soap phase,

optionally separating the neutral substances from the hydrocarbon phase,
separating sterols from the hydrocarbon phase containing neutral substances
or from the neutral substances, and
optionally purifying the sterols.

5 The above steps (a) to (c) can be carried out as explained above for the first
aspect of the present invention.

Description of the Preferred Embodiments

A preferred method for separating neutral substances from a soap
10 containing the neutral substances comprises the steps of:

- (a) providing a mixture comprising (1) the soap containing the neutral
substances, (2) water optionally containing sodium sulfate and (3) a C_1 - C_{10}
hydrocarbon solvent, and heating the mixture to a temperature of at least 140°C ,
to obtain a soap phase and a hydrocarbon phase containing the neutral
15 substances, the heating step being conducted in a closed system under pressure,
wherein the pressure in the system is at least equal to the vapor pressure of the
mixture at the temperature used in the heating step;
- (b) separating the hydrocarbon phase from the soap phase; and
- (c) optionally separating the neutral substances from the hydrocarbon
20 phase.

A preferred method for purifying sterols from a soap containing the sterols
comprises the steps of:

- (a) providing a mixture comprising (1) the soap containing the sterols,
(2) water optionally containing sodium sulfate and (3) a C_1 - C_{10} hydrocarbon
25 solvent, and heating the mixture to a temperature of at least 140°C , to obtain a
soap phase and a hydrocarbon phase containing the sterols, the heating step
being conducted in a closed system under pressure, wherein the pressure in the

system is at least equal to the vapor pressure of the mixture at the temperature used in the heating step;

- (b) separating the hydrocarbon phase and the soap phase; and
- (c) purifying the sterols from the hydrocarbon phase.

5 Preferable soaps useable as starting material in the methods of the invention are e.g. crude soaps isolated from black liquor in the sulfate cellulose production process or saponified tall oil pitch (pitch soaps) arrived at after removing e.g. by distilling tall oil from pulping soap and saponifying the remaining tall oil pitch. The use of pitch soap is especially preferable, if sterols are to be isolated
10 from the neutral substances (unsaponifiables or neutrals), because the sterol content is very high in this process flow, i.e. about 8 to 20 weight-% of the pitch dry weight.

 The term "short chain hydrocarbon" includes hydrocarbons containing 1-10 carbon atoms, more preferably 5-8 carbon atoms. Preferred short chain
15 hydrocarbon solvents in the methods are straight chain hydrocarbons (preferably pentane, hexane, heptane or octane), branched chain hydrocarbons, hydrocarbons containing cycloaliphatic structures like cyclohexane or methylcyclohexane or hydrocarbon mixtures comprising any of these compounds. Most preferred extraction solvents are hexane and/or heptane.

20 The temperature of the extracting step is preferably between 140°C and 190°C. The extraction process can technically also be performed at higher temperatures, but it is then less economic. More preferably, the temperature is between 150 and 185°C and most preferably between 165 and 185°C.

 The length of time the temperature needs to be raised to accomplish the
25 desired result is not important. It should be at least the time needed to establish the needed contact between the hydrocarbon phase and the soap phase. The time therefore depends on the degree of mixing. In most cases, the length of time

may be several seconds only.

The pressure in the extraction step depends on the temperature; the higher the temperature, the higher the pressure. Preferably the pressure is equal to the vapor pressure at the temperature of the extraction mixture in the extraction
5 step.

In the methods of the invention, sodium sulfate may optionally be added, e.g. dissolved in the water added to the soap. Also the soap itself often contains sodium sulfate. This salt is known to enhance the extraction of the neutral substances.

10 The amount of solvents useable in the methods may vary largely. However, the typical ratio of soap (in dry weight) to water (by weight) to hydrocarbon solvent (by weight) is 1 : >1 : >1. The amounts used in the methods according to the invention are preferably such that the amount of water added is minimized, thereby avoiding the problem of handling large amounts of
15 waste water and minimizing the costs of the extraction process. Instead, more hydrocarbon may be used to make the extraction more effective. The hydrocarbon is easily separated and recirculated e.g. in a closed system. The water, on the other hand, typically has to be evaporated from the soap and recirculated or purified and let into the waste water.

20 It has been found that using water in amounts higher than 3 times the amount of dry soap does not enhance the extraction of the neutral substances. A method using an elevated amount of hydrocarbon solvent is more energy sparing than a method in which a large amount of water has to be separated from the soap. Therefore, the extraction process of the invention preferably may be
25 realized by using 1-3 times as much water as the amount of soap, most preferably 2-3 times. The amount of hydrocarbon is preferably 2-6 times the amount of dry soap, more preferably 3-6 times, most preferably 4-5 times. At

these ratios a very effective extraction can be accomplished. Preferably, the weight ratios of the soap to water to hydrocarbon is 1 : 1-3 : 2-6, more preferably 1 : 2-3 : 3-6, most preferably 1: 2-3 : 4-5, to obtain a high yield of neutral substances. The extraction yield is better the more solvents are used, but
5 the use of larger amounts of solvents leads to less economic processing of the applied solvents.

The hydrocarbon phase separated during the extraction may contain small residues of fatty and resin acid salts and inorganic salts as well. These components may be harmful impurities when separating valuable components,
10 such as sterols, from the neutral fraction. It is possible to wash these impurities from the hydrocarbon phase with water. It has been noticed that the washing procedure has improved results if performed at elevated temperatures and pressure, i.e. at more or less similar conditions as the extraction has been performed, if there is any tendency that the soap residues emulsify the washing
15 water into the hydrocarbon phase and it would thereby be impossible to get the water phase separated from the hydrocarbon phase. The water wash can suitably be performed at temperatures above 80°C, most preferably above about 130°C, in a closed system. Normally, the water wash cannot be performed in open reactor systems, because the solvents evaporate too quickly to make the process
20 efficient.

This high pressure water wash can be accomplished as a batch process or more preferably continuously using e.g. a countercurrent column or mixer-settler extraction equipment.

The separated hydrocarbon phase can be reduced by partial evaporation,
25 and this partly evaporated hydrocarbon phase can be washed with water as explained above.

Neutral substances can be separated from the hydrocarbon phase by

evaporating the same to dryness.

Neutral substances can be used for isolation of e.g. sterols, wax alcohols and/or squalene (from birch based soap). When the aim is to isolate sterols from soaps, the disclosed extraction method can be followed by one or more separation and optionally performed purification steps in order to further purify the sterols from any other neutral substances.

Thus, in the second aspect of the present invention, sterols are separated from the hydrocarbon phase containing neutral substances or from the neutral substances themselves, and optionally purified to remove any other components (e.g. impurities) in a final cleaning stage.

Some preferred ways to separate and purify the sterols are explained in the following disclosure.

From the hydrocarbon phase, or preferably from the water washed hydrocarbon phase, the sterol components may be crystallized. This may be done simply by cooling the hydrocarbon phase to a suitable temperature, e.g. to about 25°C or less, followed by filtration or using some other solids/liquid separation process. Preferably the obtained sterol crystals are washed with fresh hydrocarbon. The procedure may optionally be realized by evaporating some of the hydrocarbon, optionally washing this reduced hydrocarbon phase with water, and performing the crystallization procedures from this phase. Also in this case it is preferred to wash the obtained crystals with fresh hydrocarbon. However, it has been noticed that this crystallization procedure is not optimal, because some impurities (which typically cannot be analyzed by standard gas chromatographic methods and are therefore often called "non-elutables") tend to co-crystallize with the sterols. In addition to this, the crystal forms of sterols crystallized from hydrocarbon are usually flat, soft plates which tend to be stuck together during filtration. This usually makes the crystal separation procedures

difficult.

Therefore, if small amounts of a lower (C_1 - C_6) alkanol, especially methanol, are added to the hydrocarbon solution containing the sterols and optionally other neutral substances, better crystals are obtained. Advantageously the conditions for such a crystallization procedure may be: 1 part sterols/neutral substances, 1.5-5 parts hydrocarbon and 0.03-0.5 part alkanol, and an even more preferred composition comprises 1 part sterols/neutral substances, 1.5-3.5 parts hydrocarbon and 0.03-0.35 parts alkanol. In order to increase the sterol yield it may be advantageous to also add water to the crystallization mixture. The preferred amount may vary from 0 to 1 part water per 1 part sterols/neutral substances.

If further purification of the obtained sterols is needed, this may be accomplished by recrystallization, which preferably may be performed with e.g. a similar solvent composition as used in the first crystallization, or more preferably from a lower alkanol (preferably methanol) rich solvent. Advantageously the composition of the recrystallization solution may be 1 part sterols/neutral substances, 1-3 parts hydrocarbon and 1-15 parts alkanol.

The obtained sterol crystals from the crystallization or recrystallization are preferably washed. This can be performed using any suitable solvent. Usually it is preferred to use the same solvent or mixture of solvents that are used in the isolation process, which solvents therefore already are available at the facility. Preferably the same solvent composition as used in the crystallization or recrystallization is used.

All the crystallization and/or recrystallization procedures are performed by standard methods, i.e. by heating the mixture to dissolve the sterols, if the mixture is not hot already from the preceding process steps, and then allowing the mixture to cool to crystallize the sterols. The crystals may be isolated by e.g.

filtration or centrifugation.

To minimize the loss of sterols into the solvents used in the recrystallization, the used solvent is preferably evaporated to dryness or to a reduced solution, which preferably is recirculated to the first crystallization
5 procedure for a following batch or in a continuous process.

Especially in the case of black liquor soap neutral substances, an alternative approach in separating the sterols is to use a lower alkanol (preferably methanol) rich solvent in the first crystallization step. This procedure has proven to give very pure sterols in one step, and a further advantage is the excellent filterability
10 of the sterols. The weight composition of the crystallization mixture in this case may be: 1 part sterols/neutral substances, 1.5-4 parts alkanol, 0.01-1.0 parts hydrocarbon and 0-0.05 parts water, preferably 1 part sterols/neutral substances, 1.5-4 parts alkanol, 0.01-0.3 parts hydrocarbon and 0-0.02 parts water, and most preferably 1 : 2.5 : 0.1 : 0-0.0015. In this procedure, the washing of the
15 crystallized sterols is preferably performed with a nonpolar solvent (e.g. a hydrocarbon), because the impurities stuck to the surface of the crystals are not as soluble in a polar solvent.

If the hydrocarbon phase obtained from the extraction process as such or washed with water is dried by evaporating the solvent, any known method for
20 separating the sterols from neutral substances and optionally purifying them to remove impurities may be used. A very preferred solvent mixture for crystallizing the sterols comprises methyl ethyl ketone (MEK), a lower alkanol (preferably methanol) and water. A preferred composition comprises, by weight, 50-80 parts MEK, 5-40 parts alkanol and 2-20 parts water. A more preferred
25 composition comprises 60-70 parts MEK, 20-35 parts alkanol and 5-10 parts water. The sum of these parts is preferably 100. A typical composition comprises about 70 parts MEK, about 20 parts alkanol and about 10 parts water.

The amount of MEK in the composition is preferably at most 80% by

weight, more preferably at most 75% by weight. The amount of lower alkanol is preferably at least 14% by weight, more preferably at least 17% by weight, and the amount of water is preferably at least 6% by weight, more preferably at least 8% by weight.

- 5 The following examples illustrate the invention in more detail. All the percentages given in the examples are calculated as weight - %.

Example 1

- Soap originated from *Pinus radiata* pine wood pulp was extracted in a closed reactor as batch extraction. The soap contained 15% unsaponifiab
10 calculated from the soap solids and the unsaponifiab
500 g soap (solids content 60%) was mixed with 400 g water and 900 g hydrocarbon solvent LIAV110 (a mixture of straight chain, branched chain and cycloaliphatic saturated hydrocarbons, mainly C₆-C₈ hydrocarbons, from Neste Oy) and the temperature was elevated to 150°C. After 5 min mixing (300 rpm),
15 the phases were allowed to separate for a few minutes. The lower phase was drained out slowly from the bottom valve through a Liebig type water cooled bomb. No clear intermediate ("rag") layer was detected between the lower soap layer and upper hydrocarbon layer. When evaporated to dryness, 36 g (yield 80%) unsaponifiab
20 12.3 g sterols, which was 78% of the sterols in the feed soap.

Example 2

The extraction procedure of example 1 was performed using pure heptane as a solvent. No emulsion formation was noticed. The recovery of the unsaponifiab
was 78% (recovery of sterols 77%).

Example 3

Saponified pitch, mainly originated from tall oil of *Pinus taedapine* wood pulping, was used as the material to be extracted. The solids content was 65%, the amount of unsaponifiabiles 31% and the amount of sterols 36% of the unsaponifiabiles. The extraction was performed as a single stage batch extraction as disclosed in example 1. The extraction solvent was LIAV110 and the ratios of dry pitch soap 1.0, total water 2.0 and solvent 3.9 parts. The extraction temperature was 168°C. No clear middle layer was observed. The recovery of the unsaponifiabiles and the sterols into the hydrocarbon phase was 79 and 75%, respectively.

Example 4

The same kind of soap used as raw material in example 1 was extracted continuously and countercurrently in a two-stage mixer-settler type extraction equipment. Soap, water and solvent (LIAV110) were mixed in static mixers and settling of the phases took place in settling tanks. Solvent feed was 384 kg/h, additional water feed 150 kg/h, and soap feed 131 kg/h. The temperature in both settlers was 178°C and the pressure in the first decanter was 19.2 bars, and in the second, 18.7 bars. In the interphase between the soap and the hydrocarbon layers only a small middle layer was observed. The recovery of the unsaponifiabiles into hydrocarbon in this two-stage extraction was 96% and the recovery of sterols was 94%.

Example 5

Pitch soap (*Pinus taeda* based, unsaponifiabiles 29.6%, sterols 41% of the unsaponifiabiles) was extracted with the commercial hydrocarbon mixture LIAV110 in a two-stage mixer-settler extraction equipment (the same as in

example 4). The feed ratios of dry soap to water to hydrocarbon solvent were 1:2.4:4.7 and the total feed rate was 730 kg/h. The temperatures in the settler vessels were 182°C and 184°C and the pressures were 20.7 and 20 bars, respectively. No interfering interphase layer was observed during the extraction.

- 5 The recoveries of unsaponifiabiles and sterols into the hydrocarbon phase were 95.8% and 97%, respectively.

Example 6

The extraction procedure disclosed in example 5 was performed using the feed ratios 1:1.3:6.4. The total feed rate was 570 kg/h, and the temperatures and pressures were 178°C and 19.3 bars, and 182°C and 18.9 bars, respectively.
10 The yield of unsaponifiabiles and sterols into the hydrocarbon phase was 97.4 and 97.9%, respectively.

Example 7

The starting material in this process was the unsaponifiabiles separated from tall oil pitch (100% pine origin). The pitch was first saponified at 170°C with NaOH, and then the unsaponifiabiles were extracted using a hydrocarbon solvent as in example 5 at the saponification temperature and prevailing vapor pressure in the system.
15

Example 8

20 The hydrocarbon phase (solids content 11.3%) from example 5 was washed with water as a continuous extraction process in a one-stage mixer-settler equipment. The feed rate of the hydrocarbon phase was 320 kg/h and the feed rate of the water was 100 kg/h. The temperature was 160°C. The ash content of the solids in the dried neutrals before the wash was 0.3%, and after wash,

0.04 %. According to FTIR analyses the substance transferred to the water phase consisted mainly of carboxylic acid salts.

Example 9

The hydrocarbon phase from example 8 was cooled to 20°C, and the
5 precipitated sterols were filtered and washed with fresh hydrocarbon solvent. When 100 g of dry neutrals was used as starting material, 22.5 g of a sterol blend consisting of sitosterol, sitostanol, campesterol and campestanol and nonelutable impurities was obtained. The sterol content was 80%.

Example 10

10 10 g sterols (80%) from example 9 were dissolved in a solvent consisting of 20 g LIAV110 and 90 g methanol by refluxing. The sterols were recrystallized by cooling, separated by filtration and washed with fresh recrystallization solvent. The amount of crystallized product was 6.9 g of 97% pure sterols.

15 Example 11

The hydrocarbon phase of example 8 was evaporated to a concentration where the neutrals/LIAV110 ratio was 1:2.5. 0.05 parts of MeOH was added calculated on the amount of neutrals and the blend was refluxed resulting in a clear solution. After cooling to 25°C under gentle magnetic stirring, the formed
20 crystals were filtered and washed with crystallization solvent (LIAV:MeOH 2.5:0.05). The sterol yield was 67.3% of the sterol content in the neutrals and the purity was 96.5%.

Example 12

The hydrocarbon phase of example 8 was evaporated to a concentration where the neutrals/LIAV110 ratio was 1:3.2. 0.35 parts of MeOH calculated on the amount of neutrals in the hydrocarbon phase and 1 part of water was added, and the mixture was refluxed to yield a clear solution. After crystallization, filtration and washing with a solvent of the same composition as the crystallization solvent, 73.4% of a 95.5% pure sterol blend was obtained.

Example 13

The starting material was the hydrocarbon extract from example 4 that contains unsaponifiabiles. The solvent was evaporated so that the neutrals:LIAV ratio was 1:0.1. 2.5 parts of methanol was added, calculated from the amount of neutrals, and the mixture was refluxed. The crystallization was accomplished by cooling to about 25°C, the crystals were filtered and washed with pure LIAV110. The dried product was a 97.5% pure sterol blend. The sterol recovery was 63.5% of the original sterol content in the hydrocarbon extract used as feed solution.

Example 14

The same procedure as described in example 13 was performed. The difference in this case was that the methanol used contained 1.5% water. The sterol purity was 95.8% and the yield 74.9% of the original sterol content in the hydrocarbon phase.

Example 15

The extraction solvent was distilled off from the hydrocarbon phase obtained in example 7. Then the fraction of unsaponifiabiles (35% sterols) was dissolved by refluxing into a solvent mixture consisting of 65% MEK, 30%

MeOH and 5% water (10 g unsaponifiables/ 80 g solvent). Crystallization of the sterols took place when the mixture was allowed to cool to 25°C for 1.5 h by gentle mixing with a magnetic stirrer. The crystals were filtered and washed with fresh crystallization solvent. 2.8 g of 96.5% pure dry sterol was obtained.

What is claimed is:

1. A method for separating neutral substances from a soap containing the neutral substances, the method comprising the steps of:
 - (a) providing a mixture comprising (1) the soap containing the neutral substances, (2) water optionally containing sodium sulfate and (3) a C₁-C₁₀ hydrocarbon solvent, and heating the mixture to a temperature of at least 140°C, to obtain a soap phase and a hydrocarbon phase containing the neutral substances, the heating step being conducted in a closed system under pressure, wherein the pressure in the system is at least equal to the vapor pressure of the mixture at the temperature used in the heating step;
 - (b) separating the hydrocarbon phase from the soap phase; and
 - (c) optionally separating the neutral substances from the hydrocarbon phase.
2. The method of claim 1, wherein the hydrocarbon solvent is selected from the group consisting of hexane, heptane, octane, cyclohexane, methylcyclohexane and mixtures thereof.
3. The method of claim 1, wherein the temperature is between 140°C and 190°C
4. The method of claim 3, wherein the temperature is between 165°C and 185°C.
5. The method of claim 1, wherein the soap, the water and the hydrocarbon solvent are provided in the mixture in a weight ratio of 1 : > 1 : > 1, based on

the dry weight of the soap.

6. The method of claim 5, wherein the weight ratio is 1 : 1-3 : 2-6.
7. The method of claim 5, wherein the weight ratio is 1 : 2-3 : 3-6.
8. The method of claim 5, wherein the weight ratio is 1 : 2-3 : 4-5.
- 5 9. The method of claim 1, comprising the further step of reducing the separated hydrocarbon phase produced in step (b) by evaporation.
10. The method of claim 1, comprising the further step of washing the separated hydrocarbon phase produced in step (b) with water.
11. The method of claim 9, comprising the further step of washing the reduced
10 hydrocarbon phase produced in said further step with water.
12. The method of claim 10, wherein said washing step is performed at a temperature of at least 80°C under pressure.
13. The method of claim 11, wherein said washing step is performed at a temperature of at least 80°C under pressure.
- 15 14. The method of claim 12, wherein the temperature is at least 130°C.
15. The method of claim 13, wherein the temperature is at least 130°C.

16. The method of claim 1, wherein said separating step (c) comprises evaporating the hydrocarbon phase to dryness.

17. The method of claim 1, wherein the neutral substances are selected from the group consisting of sterols, terpene alcohols and fatty alcohols.

5 18. A method for purifying sterols from a soap containing the sterols, the method comprising the steps of:

- (a) providing a mixture comprising (1) the soap containing the sterols, (2) water optionally containing sodium sulfate and (3) a C₁-C₁₀ hydrocarbon solvent, and heating the mixture to a temperature of at least 140°C, to obtain a
10 soap phase and a hydrocarbon phase containing the sterols, the heating step being conducted in a closed system under pressure, wherein the pressure in the system is at least equal to the vapor pressure of the mixture at the temperature used in the heating step;
- (b) separating the hydrocarbon phase and the soap phase; and
- 15 (c) purifying the sterols from the hydrocarbon phase.

19. The method of claim 18, wherein the hydrocarbon solvent is selected from the group consisting of hexane, heptane, octane, cyclohexane, methylcyclohexane and mixtures thereof.

20 20. The method of claim 18, wherein the temperature is between 140°C and 190°C.

21. The method of claim 20, wherein the temperature is between 165°C and 185°C.

22. The method of claim 18, wherein the soap, the water and the hydrocarbon solvent are provided in the mixture in a weight ratio of 1 : > 1 : > 1, based on the dry weight of the soap.
23. The method of claim 22, wherein the weight ratio is 1 : 1-3 : 2-6.
- 5 24. The method of claim 22, wherein the weight ratio is 1 : 2-3 : 3-6.
25. The method of claim 22, wherein the weight ratio is 1 : 2-3 : 4-5.
26. The method of claim 18, comprising the further step of reducing the separated hydrocarbon phase produced in step (b) by evaporation.
- 10 27. The method of claim 18, comprising the further step of washing the separated hydrocarbon phase produced in step (b) with water.
28. The method of claim 26, comprising the further step of washing the reduced hydrocarbon phase produced in said further step with water.
29. The method of claim 27, wherein said washing step is performed at a temperature of at least 80°C under pressure.
- 15 30. The method of claim 28, wherein said washing step is performed at a temperature of at least 80°C under pressure.
31. The method of claim 29, wherein the temperature is at least 130°C.

32. The method of claim 30, wherein the temperature is at least 130°C.

33. The method of claim 18, wherein said purifying step (c) comprises evaporating the hydrocarbon phase to dryness.

34. The method of claim 18, wherein the hydrocarbon phase additionally
5 contains other neutral substances from the soap, and step (c) further comprises purifying the sterols from the other neutral substances by dissolving the other neutral substances in a solvent mixture comprising methyl ethyl ketone, a C₁-C₆ alkanol and water, and thereafter crystallizing the sterols from the solvent mixture.

10 35. The method of claim 34, wherein the methyl ethyl ketone, the C₁-C₆ alkanol and the water are present in the solvent mixture in a weight ratio of 50-80 : 5-40 : 2-20.

36. The method of claim 35, wherein the weight ratio is 60-70 : 20-35 : 5-10.

15 37. The method of claim 34, comprising the further step of washing the sterol crystals with a solvent.

38. The method of claim 37, wherein the solvent is a mixture comprising methyl ethyl ketone, a C₁-C₆ alkanol and water.

39. The method of claim 18, wherein step (c) comprises crystallizing the
sterols from the hydrocarbon phase by cooling the hydrocarbon phase and
20 thereafter separating the formed crystals from the hydrocarbon phase.

40. The method claim 39, comprising the further step of washing the crystals and/or recrystallizing the sterols.

41. The method of claim 40, wherein the sterols are recrystallized in a solvent mixture comprising a hydrocarbon and a C₁-C₆ alkanol.

5 42. The method of claim 41, wherein the sterols, the hydrocarbon and the C₁-C₆ alkanol are present in the solvent mixture in a weight ratio of 1 : 1-3 : 1-15.

43. The method of claim 18, wherein step (c) comprises reducing the hydrocarbon phase by evaporation, mixing the reduced hydrocarbon phase with a C₁-C₆ alkanol and optionally water, and crystallizing the sterols from the
10 mixture.

44. The method of claim 43, wherein the sterols, the C₁-C₆ alkanol, the hydrocarbon and the water are present in a weight ratio of 1 : 1.5-4 : 0.01-1.0 : 0-0.05, based on the dry weight of the sterols.

45. The method of claim 44, wherein the weight ratio is 1 : 1.5-4 : 0.01-0.3
15 : 0-0.02.

46. The method claim 34, wherein the C₁-C₆ alkanol is methanol.

47. The method claim 43, wherein the C₁-C₆ alkanol is methanol.

INTERNATIONAL SEARCH REPORT

International Application No

PCT 00/00542

A. CLASSIFICATION OF SUBJECT MATTER
IPC 7 C11B13/00

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 7 C11B C11D C07J C09F

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

WPI Data, EPO-Internal, PAJ, CHEM ABS Data, PIRA

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	WO 96 10033 A (UNIV BRITISH COLUMBIA) 4 April 1996 (1996-04-04) page 9, line 16 -page 10, line 10; example 1	1-8, 17-25, 34-47
A	--- DATABASE CAPLUS 'Online! CHEMICAL ABSTRACTS; AN 1979:493367, XP002143291 abstract & FI 55 679 C (LINOTEK OY) 22 September 1978 (1978-09-22) cited in the application --- -/--	1,3-9, 17,18, 20-26,39



Further documents are listed in the continuation of box C.



Patent family members are listed in annex.

* Special categories of cited documents :

- "A" document defining the general state of the art which is not considered to be of particular relevance
- "E" earlier document but published on or after the international filing date
- "L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
- "O" document referring to an oral disclosure, use, exhibition or other means
- "P" document published prior to the international filing date but later than the priority date claimed

- "T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
- "X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
- "Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.
- "&" document member of the same patent family

Date of the actual completion of the international search

24 July 2000

Date of mailing of the international search report

07/08/2000

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2
NL - 2280 HV Rijswijk
Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,
Fax: (+31-70) 340-3016

Authorized officer

Saunders, T

INTERNATIONAL SEARCH REPORT

International Application No

PCT/IB 00/00542

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	US 3 965 085 A (HOLMBOM BJARNE ET AL) 22 June 1976 (1976-06-22) cited in the application examples 3,5	1-8, 17-26, 33-36
A	EP 0 122 322 A (UNION CAMP CORP) 24 October 1984 (1984-10-24) claims	1,3-8, 17,18, 20-25

INTERNATIONAL SEARCH REPORT

Information on patent family members

International Application No

PCT/90/00542

Patent document cited in search report		Publication date	Patent family member(s)	Publication date
WO 9610033	A	04-04-1996	AU 3559795 A	19-04-1996
			BG 101354 A	31-03-1998
			BR 9509119 A	28-10-1997
			CA 2201328 A	04-04-1996
			CN 1166174 A	26-11-1997
			CZ 9700927 A	18-02-1998
			EP 0783514 A	16-07-1997
			HU 77522 A	28-05-1998
			JP 10506394 T	23-06-1998
			LT 97074 A, B	27-10-1997
			LV 11900 A	20-12-1997
			LV 11900 B	20-05-1998
			MD 970178 A	28-02-1999
			NO 971304 A	14-05-1997
			PL 319384 A	04-08-1997
			SK 41597 A	05-11-1997
			US 5770749 A	23-06-1998
FI 55679	C	22-09-1978	FI 770873 A, B,	22-09-1978
US 3965085	A	22-06-1976	CA 1021221 A	22-11-1977
			JP 50038705 A	10-04-1975
			SE 417107 B	23-02-1981
			SE 7407901 A	30-12-1974
EP 0122322	A	24-10-1984	US 4422966 A	27-12-1983
			AT 25400 T	15-02-1987
			CA 1202302 A	25-03-1986
			FI 834818 A, B,	19-09-1984

PATENT COOPERATION TREATY

From the
INTERNATIONAL PRELIMINARY EXAMINING AUTHORITY

To:

BERMAN, Richard J.
Arent Fox Kintner Plotkin & Kahn
PLLC
1050 Connecticut Avenue, N.W.
Suite 600
Washington, DC 20036-5339
ETATS-UNIS D'AMERIQUE

PCT

NOTIFICATION OF TRANSMITTAL OF THE INTERNATIONAL PRELIMINARY EXAMINATION REPORT (PCT Rule 71.1)

Date of mailing (day/month/year)	12.03.2001
-------------------------------------	------------

Applicant's or agent's file reference 023406.00003	IMPORTANT NOTIFICATION
---	-------------------------------

International application No. PCT/IB00/00542	International filing date (day/month/year) 27/04/2000	Priority date (day/month/year) 27/04/1999
---	--	--

Applicant STEROL TECHNOLOGIES LTD. et al.
--

1. The applicant is hereby notified that this International Preliminary Examining Authority transmits herewith the international preliminary examination report and its annexes, if any, established on the international application.
2. A copy of the report and its annexes, if any, is being transmitted to the International Bureau for communication to all the elected Offices.
3. Where required by any of the elected Offices, the International Bureau will prepare an English translation of the report (but not of any annexes) and will transmit such translation to those Offices.

4. REMINDER

The applicant must enter the national phase before each elected Office by performing certain acts (filing translations and paying national fees) within 30 months from the priority date (or later in some Offices) (Article 39(1)) (see also the reminder sent by the International Bureau with Form PCT/IB/301).

Where a translation of the international application must be furnished to an elected Office, that translation must contain a translation of any annexes to the international preliminary examination report. It is the applicant's responsibility to prepare and furnish such translation directly to each elected Office concerned.

For further details on the applicable time limits and requirements of the elected Offices, see Volume II of the PCT Applicant's Guide.

Name and mailing address of the IPEA/	Authorized officer
---------------------------------------	--------------------



European Patent Office
D-80298 Munich
Tel. +49 89 2399 - 0 Tx: 523656 epmu d
Fax: +49 89 2399 - 4465

Le Bolloch, C

Tel. +49 89 2399-8091



PATENT COOPERATION TREATY

PCT

INTERNATIONAL PRELIMINARY EXAMINATION REPORT

(PCT Article 36 and Rule 70)

Applicant's or agent's file reference 023406.00003	FOR FURTHER ACTION		See Notification of Transmittal of International Preliminary Examination Report (Form PCT/IPEA/416)
International application No. PCT/IB00/00542	International filing date (day/month/year) 27/04/2000	Priority date (day/month/year) 27/04/1999	
International Patent Classification (IPC) or national classification and IPC C11B13/00			
Applicant STEROL TECHNOLOGIES LTD. et al.			

1. This international preliminary examination report has been prepared by this International Preliminary Examining Authority and is transmitted to the applicant according to Article 36.

2. This REPORT consists of a total of 5 sheets, including this cover sheet.

- ☐ This report is also accompanied by ANNEXES, i.e. sheets of the description, claims and/or drawings which have been amended and are the basis for this report and/or sheets containing rectifications made before this Authority (see Rule 70.16 and Section 607 of the Administrative Instructions under the PCT).

These annexes consist of a total of sheets.

3. This report contains indications relating to the following items:

- I ☒ Basis of the report
- II ☐ Priority
- III ☐ Non-establishment of opinion with regard to novelty, inventive step and industrial applicability
- IV ☐ Lack of unity of invention
- V ☒ Reasoned statement under Article 35(2) with regard to novelty, inventive step or industrial applicability; citations and explanations supporting such statement
- VI ☐ Certain documents cited
- VII ☐ Certain defects in the international application
- VIII ☐ Certain observations on the international application

Date of submission of the demand 27/11/2000	Date of completion of this report 12.03.2001
Name and mailing address of the international preliminary examining authority: <div style="display: flex; align-items: center;"> <div> European Patent Office D-80298 Munich Tel. +49 89 2399 - 0 Tx: 523656 epmu d Fax: +49 89 2399 - 4465 </div> </div>	Authorized officer Lippert, S Telephone No. +49 89 2399 8514



INTERNATIONAL PRELIMINARY EXAMINATION REPORT

International application No. PCT/IB00/00542

I. Basis of the report

1. This report has been drawn on the basis of *(substitute sheets which have been furnished to the receiving Office in response to an invitation under Article 14 are referred to in this report as "originally filed" and are not annexed to the report since they do not contain amendments (Rules 70.16 and 70.17).):*

Description, pages:

1-19 as originally filed

Claims, No.:

1-47 as originally filed

2. With regard to the **language**, all the elements marked above were available or furnished to this Authority in the language in which the international application was filed, unless otherwise indicated under this item.

These elements were available or furnished to this Authority in the following language: , which is:

- ☐ the language of a translation furnished for the purposes of the international search (under Rule 23.1(b)).
 - ☐ the language of publication of the international application (under Rule 48.3(b)).
 - ☐ the language of a translation furnished for the purposes of international preliminary examination (under Rule 55.2 and/or 55.3).
3. With regard to any **nucleotide and/or amino acid sequence** disclosed in the international application, the international preliminary examination was carried out on the basis of the sequence listing:
- ☐ contained in the international application in written form.
 - ☐ filed together with the international application in computer readable form.
 - ☐ furnished subsequently to this Authority in written form.
 - ☐ furnished subsequently to this Authority in computer readable form.
 - ☐ The statement that the subsequently furnished written sequence listing does not go beyond the disclosure in the international application as filed has been furnished.
 - ☐ The statement that the information recorded in computer readable form is identical to the written sequence listing has been furnished.
4. The amendments have resulted in the cancellation of:
- ☐ the description, pages:
 - ☐ the claims, Nos.:
 - ☐ the drawings, sheets:
5. ☐ This report has been established as if (some of) the amendments had not been made, since they have been considered to go beyond the disclosure as filed (Rule 70.2(c)):

INTERNATIONAL PRELIMINARY EXAMINATION REPORT

International application No. PCT/IB00/00542

(Any replacement sheet containing such amendments must be referred to under item 1 and annexed to this report.)

6. Additional observations, if necessary:

V. Reasoned statement under Article 35(2) with regard to novelty, inventive step or industrial applicability; citations and explanations supporting such statement

1. Statement

Novelty (N)	Yes: Claims 1-47
	No: Claims
Inventive step (IS)	Yes: Claims 1-47
	No: Claims
Industrial applicability (IA)	Yes: Claims 1-47
	No: Claims

2. Citations and explanations see separate sheet

ad V:

1. The claimed subject-matter is considered to be novel as none of the prior art documents cited in the International Search Report discloses all the essential features of the present application (Art.33(2) PCT).
 - 1.1 WO96/10033 (D1) discloses a process for purifying and preparing phytosterol compositions from pulping soap extracting from the pulping soap a precipitate and purifying this precipitate. For extracting the sterols a hydrocarbon solvent is used, cf. page 9, line 17 of D1 (e.g. hexane, line 28). It is also mentioned that the step can be performed at temperatures generally from 25 to 150°C, preferably between 50 and 100°C. No temperatures of at least 140°C are mentioned in the examples. It is also not explicitly mentioned that the heating step is conducted in a closed system under pressure of at least the vapor pressure of the mixture (at the temperature used in the heating step).
 - 1.2 FI-B- 55679 (D2), abstract, discloses the removal of nonsaponic matter from raw soap by heating at 100°C and normal pressure with an aqueous C₁₂₋₃₅ paraffin solvent and then heating under pressure at 160°C to give a paraffin phase containing the nonsaponic matter. No (short chain) C₁-C₁₀ solvent is disclosed in D2 as is used in present claimed process.
 - 1.3 US-A-3 965 085 (D3) discloses a method for separating neutral substances by means of extraction from an aqueous solution containing ketones by water immiscible organic solvents (e.g. gasoline, hexane, heptane cf. claim 6) to obtain a soap and an organic phase containing the neutral substances and a water phase containing the fatty acids, which phases are separated and processed further in that for instance the neutral substances is separated from the hydrocarbon phase. D3 does not disclose present heating step.
 - 1.4 EP-A-0 122 322 (D4) discloses a process for separating neutral compounds from crude tall oil soaps comprising contacting the soap with a supercritical fluid solvent for the tall oil neutral compounds whereby neutrals are extracted into said solvent to form a mixture of solvent and extract, separating the mixture from remainder of

**INTERNATIONAL PRELIMINARY
EXAMINATION REPORT - SEPARATE SHEET**

International application No. PCT/IB00/00542

the contacted soap and separating the solvent from the mixture. D4 is not directed to an extraction from an aqueous solution of the soap.

2. The closest prior art D2 differs from present claimed subject-matter in that a long chain hydrocarbon is used. The problem to be solved over D2 was to provide a process for separating neutral substances from different kind of soaps which should overcome the problem of difficult solvent regeneration, cf. page 3 of the application. The solution is to extract with a short chain solvent, which is neither disclosed nor suggested in D2. Also in combination with D3 one does not arrive at the present claimed subject-matter as D3 teaches to use two-component solvent, which also involves problems with solvent regeneration. The claimed subject-matter is therefore considered to be also inventive (Art.33(3) PCT).
3. Industrial applicability is given (Art.33(4) PCT).

PATENT COOPERATION TREATY

PCT

INTERNATIONAL SEARCH REPORT

(PCT Article 18 and Rules 43 and 44)

Applicant's or agent's file reference 023406.00003	FOR FURTHER ACTION see Notification of Transmittal of International Search Report (Form PCT/ISA/220) as well as, where applicable, item 5 below.	
International application No. PCT/IB 00/ 00542	International filing date (day/month/year) 27/04/2000	(Earliest) Priority Date (day/month/year) 27/04/1999
Applicant STEROL TECHNOLOGIES LTD. et al.		

This International Search Report has been prepared by this International Searching Authority and is transmitted to the applicant according to Article 18. A copy is being transmitted to the International Bureau.

This International Search Report consists of a total of 3 sheets.

☒ It is also accompanied by a copy of each prior art document cited in this report.

1. Basis of the report

a. With regard to the **language**, the international search was carried out on the basis of the international application in the language in which it was filed, unless otherwise indicated under this item.

☐ the international search was carried out on the basis of a translation of the international application furnished to this Authority (Rule 23.1(b)).

b. With regard to any **nucleotide and/or amino acid sequence** disclosed in the international application, the international search was carried out on the basis of the sequence listing :

☐ contained in the international application in written form.

☐ filed together with the international application in computer readable form.

☐ furnished subsequently to this Authority in written form.

☐ furnished subsequently to this Authority in computer readable form.

☐ the statement that the subsequently furnished written sequence listing does not go beyond the disclosure in the international application as filed has been furnished.

☐ the statement that the information recorded in computer readable form is identical to the written sequence listing has been furnished

2. ☐ **Certain claims were found unsearchable** (See Box I).

3. ☐ **Unity of invention is lacking** (see Box II).

4. With regard to the **title**,

☒ the text is approved as submitted by the applicant.

☐ the text has been established by this Authority to read as follows:

5. With regard to the **abstract**,

☒ the text is approved as submitted by the applicant.

☐ the text has been established, according to Rule 38.2(b), by this Authority as it appears in Box III. The applicant may, within one month from the date of mailing of this international search report, submit comments to this Authority.

6. The figure of the **drawings** to be published with the abstract is Figure No.

☐ as suggested by the applicant.

☐ because the applicant failed to suggest a figure.

☐ because this figure better characterizes the invention.

☐ None of the figures.

INTERNATIONAL SEARCH REPORT

International Application No

IB 00/00542

A. CLASSIFICATION OF SUBJECT MATTER
IPC 7 C11B13/00

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 7 C11B C11D C07J C09F

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

WPI Data, EPO-Internal, PAJ, CHEM ABS Data, PIRA

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	WO 96 10033 A (UNIV BRITISH COLUMBIA) 4 April 1996 (1996-04-04) page 9, line 16 -page 10, line 10; example 1	1-8, 17-25, 34-47
A	--- DATABASE CAPLUS 'Online! CHEMICAL ABSTRACTS; AN 1979:493367, XP002143291 abstract & FI 55 679 C (LINOTEK OY) 22 September 1978 (1978-09-22) cited in the application --- -/--	1,3-9, 17,18, 20-26,39

☒ Further documents are listed in the continuation of box C.

☒ Patent family members are listed in annex.

* Special categories of cited documents:

"A" document defining the general state of the art which is not considered to be of particular relevance

"E" earlier document but published on or after the international filing date

"L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)

"O" document referring to an oral disclosure, use, exhibition or other means

"P" document published prior to the international filing date but later than the priority date claimed

"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

"X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

"Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.

"&" document member of the same patent family

Date of the actual completion of the international search

24 July 2000

Date of mailing of the international search report

07/08/2000

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2
NL - 2280 HV Rijswijk
Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,
Fax: (+31-70) 340-3016

Authorized officer

Saunders, T

INTERNATIONAL SEARCH REPORT

International Application No

IB 00/00542

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category °	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	US 3 965 085 A (HOLMBOM BJARNE ET AL) 22 June 1976 (1976-06-22) cited in the application examples 3,5 ---	1-8, 17-26, 33-36
A	EP 0 122 322 A (UNION CAMP CORP) 24 October 1984 (1984-10-24) claims -----	1,3-8, 17,18, 20-25

INTERNATIONAL SEARCH REPORT

Information on patent family members

International Application No

IB 00/00542

Patent document cited in search report		Publication date	Patent family member(s)	Publication date
WO 9610033	A	04-04-1996	AU 3559795 A	19-04-1996
			BG 101354 A	31-03-1998
			BR 9509119 A	28-10-1997
			CA 2201328 A	04-04-1996
			CN 1166174 A	26-11-1997
			CZ 9700927 A	18-02-1998
			EP 0783514 A	16-07-1997
			HU 77522 A	28-05-1998
			JP 10506394 T	23-06-1998
			LT 97074 A, B	27-10-1997
			LV 11900 A	20-12-1997
			LV 11900 B	20-05-1998
			MD 970178 A	28-02-1999
			NO 971304 A	14-05-1997
			PL 319384 A	04-08-1997
			SK 41597 A	05-11-1997
			US 5770749 A	23-06-1998
FI 55679	C	22-09-1978	FI 770873 A, B,	22-09-1978
US 3965085	A	22-06-1976	CA 1021221 A	22-11-1977
			JP 50038705 A	10-04-1975
			SE 417107 B	23-02-1981
			SE 7407901 A	30-12-1974
EP 0122322	A	24-10-1984	US 4422966 A	27-12-1983
			AT 25400 T	15-02-1987
			CA 1202302 A	25-03-1986
			FI 834818 A, B,	19-09-1984



INTERNATIONAL PRELIMINARY EXAMINATION REPORT

(PCT Article 36 and Rule 70)

Applicant's or agent's file reference 023406.00003	FOR FURTHER ACTION See Notification of Transmittal of International Preliminary Examination Report (Form PCT/IPEA/416)	
International application No. PCT/IB00/00542	International filing date (day/month/year) 27/04/2000	Priority date (day/month/year) 27/04/1999
International Patent Classification (IPC) or national classification and IPC C11B13/00		
Applicant STEROL TECHNOLOGIES LTD. et al.		

1. This international preliminary examination report has been prepared by this International Preliminary Examining Authority and is transmitted to the applicant according to Article 36.



2. This REPORT consists of a total of 5 sheets, including this cover sheet.

- ☐ This report is also accompanied by ANNEXES, i.e. sheets of the description, claims and/or drawings which have been amended and are the basis for this report and/or sheets containing rectifications made before this Authority (see Rule 70.16 and Section 607 of the Administrative Instructions under the PCT).

These annexes consist of a total of sheets.

3. This report contains indications relating to the following items:

- I ☒ Basis of the report
- II ☐ Priority
- III ☐ Non-establishment of opinion with regard to novelty, inventive step and industrial applicability
- IV ☐ Lack of unity of invention
- V ☒ Reasoned statement under Article 35(2) with regard to novelty, inventive step or industrial applicability; citations and explanations supporting such statement
- VI ☐ Certain documents cited
- VII ☐ Certain defects in the international application
- VIII ☐ Certain observations on the international application

Date of submission of the demand 27/11/2000	Date of completion of this report 12.03.2001
Name and mailing address of the international preliminary examining authority:  European Patent Office D-80298 Munich Tel. +49 89 2399 - 0 Tx: 523656 epmu d Fax: +49 89 2399 - 4465	Authorized officer Lippert, S Telephone No. +49 89 2399 8514 

INTERNATIONAL PRELIMINARY EXAMINATION REPORT

International application No. PCT/IB00/00542

I. Basis of the report

1. This report has been drawn on the basis of *(substitute sheets which have been furnished to the receiving Office in response to an invitation under Article 14 are referred to in this report as "originally filed" and are not annexed to the report since they do not contain amendments (Rules 70.16 and 70.17).):*

Description, pages:

1-19 as originally filed

Claims, No.:

1-47 as originally filed

2. With regard to the **language**, all the elements marked above were available or furnished to this Authority in the language in which the international application was filed, unless otherwise indicated under this item.

These elements were available or furnished to this Authority in the following language: , which is:

- ☐ the language of a translation furnished for the purposes of the international search (under Rule 23.1(b)).
- ☐ the language of publication of the international application (under Rule 48.3(b)).
- ☐ the language of a translation furnished for the purposes of international preliminary examination (under Rule 55.2 and/or 55.3).

3. With regard to any **nucleotide and/or amino acid sequence** disclosed in the international application, the international preliminary examination was carried out on the basis of the sequence listing:

- ☐ contained in the international application in written form.
- ☐ filed together with the international application in computer readable form.
- ☐ furnished subsequently to this Authority in written form.
- ☐ furnished subsequently to this Authority in computer readable form.
- ☐ The statement that the subsequently furnished written sequence listing does not go beyond the disclosure in the international application as filed has been furnished.
- ☐ The statement that the information recorded in computer readable form is identical to the written sequence listing has been furnished.

4. The amendments have resulted in the cancellation of:

- ☐ the description, pages:
- ☐ the claims, Nos.:
- ☐ the drawings, sheets:

5. ☐ This report has been established as if (some of) the amendments had not been made, since they have been considered to go beyond the disclosure as filed (Rule 70.2(c)):

INTERNATIONAL PRELIMINARY EXAMINATION REPORT

International application No. PCT/IB00/00542

(Any replacement sheet containing such amendments must be referred to under item 1 and annexed to this report.)

6. Additional observations, if necessary:

V. Reasoned statement under Article 35(2) with regard to novelty, inventive step or industrial applicability; citations and explanations supporting such statement

1. Statement

Novelty (N)	Yes:	Claims	1-47
	No:	Claims	
Inventive step (IS)	Yes:	Claims	1-47
	No:	Claims	
Industrial applicability (IA)	Yes:	Claims	1-47
	No:	Claims	

2. Citations and explanations see separate sheet

ad V:

1. The claimed subject-matter is considered to be novel as none of the prior art documents cited in the International Search Report discloses all the essential features of the present application (Art.33(2) PCT).
 - 1.1 WO96/10033 (D1) discloses a process for purifying and preparing phytosterol compositions from pulping soap extracting from the pulping soap a precipitate and purifying this precipitate. For extracting the sterols a hydrocarbon solvent is used, cf. page 9, line 17 of D1 (e.g. hexane, line 28). It is also mentioned that the step can be performed at temperatures generally from 25 to 150°C, preferably between 50 and 100°C. No temperatures of at least 140°C are mentioned in the examples. It is also not explicitly mentioned that the heating step is conducted in a closed system under pressure of at least the vapor pressure of the mixture (at the temperature used in the heating step).
 - 1.2 FI-B- 55679 (D2), abstract, discloses the removal of nonsaponic matter from raw soap by heating at 100°C and normal pressure with an aqueous C₁₂₋₃₅ paraffin solvent and then heating under pressure at 160°C to give a paraffin phase containing the nonsaponds. No (short chain) C₁-C₁₀ solvent is disclosed in D2 as is used in present claimed process.
 - 1.3 US-A-3 965 085 (D3) discloses a method for separating neutral substances by means of extraction from an aqueous solution containing ketones by water immiscible organic solvents (e.g. gasoline, hexane, heptane cf. claim 6) to obtain a soap and an organic phase containing the neutral substances and a water phase containing the fatty acids, which phases are separated and processed further in that for instance the neutral substances is separated from the hydrocarbon phase. D3 does not disclose present heating step.
 - 1.4 EP-A-0 122 322 (D4) discloses a process for separating neutral compounds from crude tall oil soaps comprising contacting the soap with a supercritical fluid solvent for the tall oil neutral compounds whereby neutrals are extracted into said solvent to form a mixture of solvent and extract, separating the mixture from remainder of

**INTERNATIONAL PRELIMINARY
EXAMINATION REPORT - SEPARATE SHEET**

International application No. PCT/IB00/00542

the contacted soap and separating the solvent from the mixture. D4 is not directed to an extraction from an aqueous solution of the soap.

2. The closest prior art D2 differs from present claimed subject-matter in that a long chain hydrocarbon is used. The problem to be solved over D2 was to provide a process for separating neutral substances from different kind of soaps which should overcome the problem of difficult solvent regeneration, cf. page 3 of the application. The solution is to extract with a short chain solvent, which is neither disclosed nor suggested in D2. Also in combination with D3 one does not arrive at the present claimed subject-matter as D3 teaches to use two-component solvent, which also involves problems with solvent regeneration. The claimed subject-matter is therefore considered to be also inventive (Art.33(3) PCT).
3. Industrial applicability is given (Art.33(4) PCT).